



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

$$\therefore \sum_{n=1}^{\infty} \frac{(n+2)^2}{n(n+4)} = n \left[ 1 + \frac{1}{n+1} + \frac{1}{2(n+2)} + \frac{1}{3(n+3)} + \frac{1}{4(n+4)} \right].$$

Also solved by Professor G. B. M. ZERR.

35. Proposed by COOPER D. SCHMITT, A. M., Professor of Mathematics, University of Tennessee, Knoxville, Tennessee.

Prove that the product of two numbers, each the sum of four (4) squares may be expressed as the sum of four squares in 48 different ways and unite some or all of the 48 ways.

Solution by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.

Let  $(a^2 + b^2 + c^2 + d^2)(e^2 + f^2 + g^2 + h^2) = A$  = the product of the two numbers. From Euler's Theorem we get

$$\begin{aligned} A &= (ae + bf - cg - dh)^2 + (af - be - ch + dg)^2 \\ &\quad + (-ag + bh - ce + df)^2 + (ah + bg + cf + de)^2, \\ &= (-ae + bf - cg + dh)^2 + (af + be + ch + dg)^2 \\ &\quad + (ag + bh - ce - df)^2 + (ah - bg - cf + de)^2, \\ &= (ae - bf - cg + dh)^2 + (af + be - ch - dg)^2 \\ &\quad + (ag + bh + ce + df)^2 + (-ah + bg - cf + de)^2, \\ &= (ae + bf + cg + dh)^2 + (-af + be - ch + dg)^2 \\ &\quad + (ag - bh - ce + df)^2 + (ah + bg - cf - de)^2, \\ &= (ae - bf + cg + dh)^2 + (af + be - ch + dg)^2 \\ &\quad + (ag - bh + ce + df)^2 + (ah + bg + cf - de)^2, \\ &= (ae + bf - cg - dh)^2 + (af + be + ch - dg)^2 \\ &\quad + (ag - bh + ce + df)^2 + (ah + bg - cf + de)^2, \\ &= (ae + bf - cg + dh)^2 + (af - be + ch + dg)^2 \\ &\quad + (ag + bh + ce - df)^2 + (ah - bg - cf - de)^2, \\ &= (ae + bf + cg - dh)^2 + (af - be - ch - dg)^2 \\ &\quad + (ag + bh - ce + df)^2 + (ah - bg + cf + de)^2. \end{aligned}$$

The sum of four squares in eight different ways by combination of signs.

$$\begin{aligned} A &= (ae + bf - cg - dh)^2 + (af - be - ch + dg)^2 \\ &\quad + (-ag + bh - ce + df)^2 + (ah + bg + cf + de)^2, \\ &= (ag + bh - ce - df)^2 + (af - bg - ce + dh)^2 \\ &\quad + (-ah + be - cg + df)^2 + (ae + bh + cf + dg)^2, \end{aligned}$$

$$\begin{aligned}
&= (ah + bf - ce - dg)^2 + (af - bh - cg + de)^2 \\
&\quad + (-ae + bg - ch + df)^2 + (ag + be + cf + dh)^2, \\
&= (ae + bg - cf - dh)^2 + (ag - be - ch + df)^2 \\
&\quad + (-af + bh - ce + dg)^2 + (ah + bf + cg + de)^2, \\
&= (af + bg - ch - de)^2 + (ag - bf - ce + dh)^2 \\
&\quad + (-ah + be - cf + dg)^2 + (ae + bh + cg + df)^2, \\
&= (ah + bg - ce - df)^2 + (ag - bh - cf + de)^2 \\
&\quad + (-af + be - cg + dh)^2 + (ae + hf + ch + dg)^2,
\end{aligned}$$

the sum of four squares in six different ways by combination of letters.

Since the signs of each of these six can form the sum of four squares in eight different ways, the whole number of ways is  $8 \times 6 = 48$  different ways.

---

## PROBLEMS.

---

46. Proposed by Professor WILLIAM HOOVER, A. M., Ph. D., Professor of Mathematics in the Ohio University, Athens, Ohio.

Find  $\theta$  from  $\cos \theta + \cos 3\theta + \cos 5\theta = 0$ .

47. Proposed by LEONARD E. DICKSON, A. M., Fellow in Mathematics, University of Chicago, Chicago, Illinois.

Prove that  $(-1)(-1) = +1$ .

---

## GEOMETRY.

---

Conducted by B. F. FINKEL, Kidder, Mo. All contributions to this department should be sent to him.

---

## SOLUTIONS OF PROBLEMS.

---

34. Proposed by T. JOHN COLE, Columbus, Ohio.

A circular field contains 10 acres. A horse is tied to the fence with a rope sufficiently long to graze over one acre. Find length of the rope (1) when the horse is on the inside (2) when he is on the outside of the fence.

Solution by Professor G. B. M. ZERR, A. M., Principal of High School, Staunton, Virginia.